

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

## **IMAGES ARE BEST AVAILABLE COPY.**

As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.

**THIS PAGE BLANK (USPTO)**

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b> <b>C07K 7/20, A61K 47/24, 37/43</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 94/28015</b> <b>(43) International Publication Date:</b> 8 December 1994 (08.12.94)
<b>(21) International Application Number:</b> PCT/AU94/00262 <b>(22) International Filing Date:</b> 20 May 1994 (20.05.94) <b>(30) Priority Data:</b> PL 8880 20 May 1993 (20.05.93) AU  <b>(71) Applicant (for all designated States except US):</b> BIOTECH AUSTRALIA PTY. LIMITED [AU/AU]; 28 Barcoo Street, Roseville, NSW 2069 (AU). <b>(72) Inventor; and</b> <b>(75) Inventor/Applicant (for US only):</b> RUSSELL-JONES, Gregory, John [AU/AU]; 23 Greenfield Avenue, Middle Cove, NSW 2068 (AU).  <b>(74) Agent:</b> F.B. RICE & CO.; 28 A Montague Street, Balmain, NSW 2041 (AU).		<b>(81) Designated States:</b> AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KG, KP, KR, KZ, LK, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> LHRH ANTAGONISTS  <b>(57) Abstract</b>  The present invention provides novel LHRH antagonists and complexes between these antagonists and VB12. The present invention is particularly useful for the oral administration of LHRH antagonists. The antagonists of the invention are selected from the group consisting of ANTIDE-1, ANTIDE-2 and ANTIDE-3.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

## LHRH ANTAGONISTS

### FIELD OF THE INVENTION

The present invention relates to LHRH antagonists and to the systemic and oral delivery of the antagonists by administration of a complex comprising the antagonists linked to vitamin B12 (VB12) or an analogue thereof. More particularly, the invention relates to methods for the synthesis of these complexes.

### 10 BACKGROUND OF THE INVENTION

Hypothalamic gonadotrophin releasing hormone (GnRH, also known as luteinizing hormone releasing hormone, LHRH) mediates the regulation of pituitary gonadotrophin synthesis and secretion. Since the initial isolation of LHRH by Schally et al. (1971) numerous analogues have been synthesized, which may be classified according to their acute effects on gonadotrophin release, as either agonists (with enhancement of release) or antagonists, (with inhibition of release). The agonists are characterised by an increase in binding affinity to pituitary LHRH receptors (Loumay et al, 1982, Perrin et al. 1980) and by increased resistance to the proteolytic degradation that rapidly removes native LHRH.

25 The observation that these potent analogues can induce potentially reversible medical castration has provided a new approach to the treatment of various gonadotrophin dependent disorders, particularly hormone dependent prostate and breast cancer. Two distinct phases in the induction of castration occur. Initially the analogue stimulates the pituitary-gonadal axis causing a transient increase in gonadotrophin and sex steroid secretion in the first two weeks or so. After this period there is a down regulation of pituitary LHRH receptors with subsequent decline in gonadotrophin and sex steroid secretion. The initial stimulation is a substantial drawback to the use of agonists in the

treatment of prostatic malignancy as the initial stimulation of testosterone can produce a painful flare of the disease with consequent adverse clinical effects (Eisenberger and Abrams, 1988, Crawford and Davis, 1988). By contrast the first successful synthesis of LHRH antagonists should be advantageous in avoiding the initial flare. First attempts at the production of antagonists led to compounds characterised by undesirable histamine release (Karten and Rivier, 1986, Schmidt et al., 1984, Phillips et al., 1988). However new antagonists have been developed characterised by improved potency and much less histamine releasing potential eg. Karten and Rivier, 1986. One of the most potent antagonists described to date is the analogue N-Ac-D-Nal(2), D-Phe (pCl), D-Pal(3), Ser, Lys (Nic), D-Lys(Nic), Leu, Lys(iPr), Pro, D-Ala-NH<sub>2</sub> (ANTIDE,) synthesized by Folkers and Bowers. It is highly potent in rats in inhibiting ovulation (Ljungquist et al., 1988) and single doses have profound, long lasting inhibitory effects on serum LH concentrations in castrate female cynomolgous monkeys (Leal et al., 1988). It has also been shown to be capable of inducing long term chemical castration in intact adult male rats and cynomolgous monkeys, and to have an inhibitory effect on tumour growth in the Dunning R3327 prostatic carcinoma model, similar to that of castration (Habenicht et al., 1990). The new antagonist should therefore be of substantial clinical interest in all those conditions in which medical castration is desirable, particularly in the management of prostatic cancer and in various gynaecological disorders (McLachlan et al., 1986). Current administration is limited to the parenteral route. However the dose of analogue that can be delivered by this route is limited due to the poor solubility of the antagonist. Thus clinical trials of the long term effect of ANTIDE have been reduced to doses of 2.5 mg or lower.

The oral route of administration of peptides such as LHRH and its analogues as pharmaceuticals in the treatment of systemic conditions has met with little success. In general the amount of peptide required for successful oral administration has been 100 to 1000 times the dose required for parenteral deliver, thus making the administration of these agents via this route prohibitively expensive. There are two fundamental reasons for the lack of success. Firstly, the intestinal milieu has a high degree of proteolytic activity, which rapidly degrades most peptides. Secondly, while there are well defined uptake mechanisms for individual amino acids and di-peptides, there is no general mechanism for polypeptides to be transported across the membrane of the mucosal epithelium into the circulation. Rather, this membrane is designed as a general barrier prohibiting the uptake of the numerous foreign proteins encountered in this environment. Thus, although a peptide may be modified to withstand the enzymatic barrage encountered in the intestine, such modification is of little value if the peptide cannot subsequently cross the mucosal barrier and enter the systemic circulation.

Recent work by the current inventor (PCT/AU86/00299) has however provided a method to overcome the mucosal barrier. This method takes advantage of the natural intrinsic factor (IF) mediated uptake mechanism for vitamin B12 (VB12). VB12 is a naturally occurring dietary molecule which is actively taken up from the intestine. During this process it first binds to IF in the upper small intestine. The VB12-IF complex then passes down the small intestine and binds to an IF receptor located on the surface of the ileal epithelium. The whole [VB12-IF-Receptor] complex is then internalized by receptor-mediated endocytosis and some time later the VB12 appears in serum.

PCT application PCT/AU86/00299 (WO87/02251) describes methods to modify chemically VB12 to provide suitable functional groups for conjugation of the VB12 to various drugs and peptide/protein pharmaceuticals.

- 5 When the [VB12-pharmaceutical] complex is administered orally it is possible to utilise the natural IF-mediated VB12-uptake system to deliver the pharmaceutical to the circulation.

- As the oral route of administration is the most  
10 preferable means of delivering a pharmaceutically active agent, there is a large and valuable market for any process which permits the oral delivery of LHRH antagonists to humans. Such a process would be available by the formation of a complex between VB12 or  
15 analogues thereof and LHRH antagonists.

#### SUMMARY OF THE INVENTION

In a first aspect the present invention consists in analogues of ANTIDE.

- ANTIDE is a particularly suitable molecule for  
20 oral administration, as it contains very few natural amino acids and as such is highly resistant to the action of pepsin, trypsin or chymotrypsin. However, ANTIDE itself needs to be modified slightly to provide a suitable site for conjugation to VB12. The inventor  
25 has found that it is possible to synthesize three analogues of ANTIDE which are suitable for conjugation to VB12, namely N-Ac-D-Nal(2), D-Phe (pCl), D-Pal(3), Ser, Lys (Nic), D-Lys, Leu, Lys(iPr), Pro, D-Ala-NH<sub>2</sub> (D-Lys<sub>6</sub>ANTIDE or ANTIDE-1), N-Ac-D-Nal(2), D-Phe (pCl),  
30 D-Pal(3), Ser, Lys, D-Lys(Nic), Leu, Lys(iPr), Pro, D-Ala-NH<sub>2</sub> (Lys<sub>5</sub> ANTIDE or ANTIDE-2) and N-Ac-D-Nal(2), D-Phe (pCl), D-Pal(3), Ser, Lys (Nic), D-Lys(Nic), Leu, Lys, Pro, D-Ala-NH<sub>2</sub> (Lys<sub>8</sub>ANTIDE or ANTIDE-3).

- A second aspect of the present invention provides  
35 a complex between the VB12 or an analogue of VB12 and the LHRH antagonists of the first aspect, wherein the ability of VB12 or an analogue of VB12 to undergo the



binding reactions necessary for the uptake and transport of VB12 in a vertebrate host and the activity of the LHRH antagonist are substantially maintained.

The VB12 analogues include any variation or derivative of VB12 which posses binding activity to intrinsic factor. Preferred analogues of VB12 include cyanocobalamin (CN-Cbl), aquocobalamin, adenosylcobalamin, methylcobalamin, hydroxycobalamin, cyanocobalamin carbanalide, 5-methoxybenzylcobalamin [(5-MeOBz)CN-Cbl], as well as the desdimethyl, monoethylamide and methylamide analogues of all of the above. Other analogues include coenzyme B12, 5'-deoxyadenosylcobalamin, chlorocobalamin, sulfitocobalamin, nitrocobalamin, thiocyanatocobalamin, benzimidazole derivatives such as 5,6-dichlorobenzimidazole, 5-hydroxybenzimidazole, trimethylbenzimidazole, as well as adenosylcyanocobalamin [(Ade)CN-Cbl], cobalamin lactone, cobalamin lactam, and the anilide, ethylamide, monocarboxylic and dicarboxylic acid derivatives of VB12 or its analogues.

Preferred derivatives of VB12 also include the mono-, di- and tricarboxylic acid derivatives of propionamide derivatives of VB12. Other derivatives include analogues of VB12 in which the cobalt is replaced by zinc or nickel. The corrin ring of VB12 or its analogues may also be substituted with any substituent which does not effect its binding to IF, and such derivatives of VB12 or its analogues are part of this invention. Other derivatives and analogues of VB12 are discussed in Schneider, Z and Stroinski, A.: (Walter De Gruyter, Berlin, NY; 1987), the disclosure of which is incorporated herein by reference.

Suitable extended spacers for the conjugation include disuccinimidyl suberate (DSS), bis(sulfosuccinimidyl) suberate (BSS), ethylene glycolbis(succinimidylsuccinate) (EGS), ethylene

glycolbis(sulfosuccinimidylsuccinate) (Sulfo-EGS), p-aminophenylacetic acid, dithiobis(succinimidylpropionate) (DSP), 3,3'-dithiobis(sulfosuccinimidylpropionate) (DTSSP),  
5 disuccinimidyl tartarate (DST), disulfosuccinimidyl tartarate (Sulfo-DST), bis[2-(succinimidooxycarbonyloxy)-ethylene]sulfone (BSOCOES), bis[2-(sulfosuccinimidooxycarbonyloxy)-ethylene]sulfone (Sulfo-BSOCOES), dimethyl adipimide.2HCl (DMA),  
10 dimethyl pimelimide.2HCl (DMP), dimethyl suberimide.2HCl (DMS).

Suitable cross-linking agents for use in the preparation of thiol-cleavable biodegradable linkers include N-succinimidyl 3-(2-pyridyldithio)propionate  
15 (SPDP), iminothiolane, sulfosuccinimidyl 6-[3-(2-pyridyldithio) propionamido] hexanoate (Sulfo-LC-SPDP), succinimidyl 6-[3-(2-pyridyldithio) propionamido] hexanoate (LC-SPDP), sulfosuccinimidyl 6-[ $\alpha$ -methyl- $\alpha$ -(2-pyridyldithio) toluamido]hexanoate (Sulfo-LC-SMPT),  
20 1,4-di[3'-(2'-pyridyldithio)propionamido]butane (DPDPB), 4-succinimidylloxycarbonyl- $\alpha$ -methyl- $\alpha$ -(2-pyridyldithio)-toluene (SMPT), dimethyl 3,3'-dithiobispropionimide.2 HCl (DTBP).

In a further aspect the invention provides a  
25 method for preparing biodegradable complexes between VB12 and LHRH antagonists such as the ANTIDE analogues of the first aspect, such that the VB12 is cleaved from the LHRH antagonist *in vivo*.

In yet another aspect the invention provides a  
30 complex comprising an ANTIDE analogue of the first aspect bound to VB12 carrier or an analogue thereof where more than one ANTIDE analogue is bound to the carrier.

In yet another aspect the invention provides a  
35 method for the oral delivery of LHRH antagonists whereby the LHRH antagonist will be transepithelially

transported from the intestine to the circulation of the vertebrate host.

In another aspect, the invention describes a new class of ANTIDE analogues which possess greater  
5 solubility in physiological solvents. These analogues can be administered systemically.

#### BRIEF DESCRIPTION OF THE DRAWINGS

In order that the nature of the present invention may be more clearly understood, preferred forms thereof  
10 will now be described with reference to the following examples and figures in which:

Figure 1 is a graph showing the median inhibitory concentrations (IC<sub>50</sub>), determined by *in vitro* bioassay, for blockade of LHRH-stimulated release of LH from rat  
15 pituitary cells by ANTIDE, its analogues and VB12-conjugates. Median inhibitory concentrations of ANTIDE, ANTIDE analogues and VB12-ANTIDE conjugates for antagonising LHRH-stimulated release of LH from rat anterior pituitary cell cultures during 4 hours.  
20 Results for conjugates are given as ng of the incorporated analogue/ml (final concentration in culture well). Results that do not differ from each other (i.e.  $P > 0.05$ ) are grouped under a common letter.

Figure 2 is a graph showing *in vitro* bioassay for  
25 blockade of LHRH-stimulated release of LH from rat pituitary cells by ANTIDE, ANTIDE-3 its VB12-conjugates. Assay conditions are similar as those given in Figure 1.

Figure 3 is a graph showing reduction in serum LH  
30 level in castrate rats following injection of ANTIDE, ANTIDE-1 or ANTIDE-1-VB12-conjugates. The *in vivo* effect of subcutaneous injection of ANTIDE or VB12-ANTIDE conjugates was examined in 6 week old castrate rats. Twenty-four hours after subcutaneous injection of  
35 a 100 µg dose of the various ANTIDE analogues, the rats were killed and trunk blood collected for analysis of serum LH levels by RIA (see Puente & Catt, 1986).

Figure 4 is a graph showing reduction in serum LH levels in castrate rats following subcutaneous injection of 100 µg doses of ANTIDE, ANTIDE-3 or ANTIDE-3-VB12-conjugates.

## 5 Materials

### Pituitary cell assay for LH release

Cell cultures were prepared from the anterior pituitaries of 4 month old Sprague-Dawley rats. Cells were dispersed as a single cell suspension of 250,000  
10 viable cells per ml in Dulbecco's modified Eagle's Medium with 10% foetal bovine serum plus antibiotics. Aliquots of 0.5 ml were dispensed into multi well tissue culture plates and maintained at 30° C in an enclosed humidified atmosphere of 5% CO<sub>2</sub> in air. Cells  
15 were cultured for 5 days and medium removed. The medium was replaced with fresh medium containing a range of doses of ANTIDE, ANTIDE analogues or VB12-ANTIDE conjugates. After 4 hours 10 nM LHRH was added to each well. Cells were incubated for a further 4 hours  
20 afterwhich medium was removed for the determination of the amount of FSH and LH released into the medium by radio-immuno assay (RIA).

### Castrate rat model

Adult Sprague Dawley rats were castrated and  
25 maintained in the animal house for one week prior to administration of test substances. ANTIDE or analogues were tested by subcutaneous injection into the castrate rats. Groups of 5 rats were used as controls and for collection of blood samples from the external jugular  
30 vein at 24 hours post-dose . LH was measured by a standard radio-immunoassay.

### IF Assay

The affinity of the various VB12-ANTIDE conjugates for IF was determined in a competitive  
35 binding assay (Russell-Jones, 1994). Dilutions of unlabelled VB12 or VB12 analogue or conjugate were mixed with 1 ng <sup>57</sup>CoVB12 (Amersham). One IU of IF (a

unit of IF is the quantity of IF required to bind 1 ng of VB12) was then added to the mixture, and the mixture was incubated for 20 min at room temperature (RT) before the addition of a solution of 5% activated charcoal in 0.1% BSA (IF and VB12-free, Sigma). Samples were centrifuged and the counts in the supernatant (IF bound) and pellet (free  $^{57}\text{CoVB12}$ ) were used to determine the relative affinity of IF for the material tested.

#### 10 Synthesis of ANTIDE analogues

Three ANTIDE analogues containing single, unmodified lysines, namely D-Lys<sup>6</sup>ANTIDE or ANTIDE-1; Lys<sup>5</sup>ANTIDE or ANTIDE-2 and Lys<sup>8</sup>ANTIDE or ANTIDE-3 (see Table 1.) were synthesized on an Applied Biosystems Peptide Synthesizer. Peptides were cleaved from the resin and purified by Reverse Phase HPLC, using a gradient of 5-100% acetonitrile in 0.1% TFA.

Table 1. Sequence of ANTIDE and its analogues.

Analogue	Sequence
ANTIDE	N-Ac-D-Nal(2), D-Phe (pCl), D-Pal(3), Ser, Lys (Nic), D-Lys(Nic), Leu, Lys(iPr), Pro, D-Ala-NH <sub>2</sub>
D-Lys <sup>6</sup> ANTIDE or ANTIDE-1	N-Ac-D-Nal(2), D-Phe (pCl), D-Pal(3), Ser, Lys (Nic), D-Lys, Leu, Lys(iPr), Pro, D-Ala-NH <sub>2</sub>
Lys <sup>5</sup> ANTIDE or ANTIDE-2	N-Ac-D-Nal(2), D-Phe (pCl), D-Pal(3), Ser, Lys, D-Lys(Nic), Leu, Lys(iPr), Pro, D-Ala-NH <sub>2</sub>
Lys <sup>8</sup> ANTIDE or ANTIDE-3	N-Ac-D-Nal(2), D-Phe (pCl), D-Pal(3), Ser, Lys (Nic), D-Lys(Nic), Leu, Lys, Pro, D-Ala-NH <sub>2</sub>

Two of the analogues, ANTIDE-1 and ANTIDE-3, have considerable potency *in vitro* and ANTIDE-3 has activity comparable to the native ANTIDE *in vivo*. The direct covalent conjugation of the "e" isomer carboxylate of VB12 (eVB12) to either ANTIDE-1, ANTIDE-2 or ANTIDE-3 was found to reduce the bioactivity of the ANTIDE analogue to levels substantially lower than ANTIDE

itself regardless of whether the analogue was tested *in vitro* or *in vivo*.

#### Direct conjugation of ANTIDE analogues to monocarboxy-VB12

5           In order to obtain a functional group suitable for conjugation of the VB12 to ANTIDE, cyanocobalamin (VB12) was first hydrolysed in 0.4 M HCl for 72 hours. The e isomer of monocarboxy vitamin B12 (eVB12, e isomer), which has previously been shown to have the  
10           highest affinity for IF of the three isomers, was separated from the b and d isomers (also formed during acid hydrolysis of cyanocobalamin) by a combination of Dowex 1X2 chromatography, and semi-preparative C-18 RP-HPLC developed with a gradient of 5-100% acetonitrile  
15           in 0.1% TFA.

          Aminoethyl-eVB12 was prepared by reacting the eVB12 isomer with 1,2-diaminoethane at pH 6.5 using a twenty-fold molar excess of the diamine over e isomer and a twenty fold molar excess of 1-ethyl-3-(3-  
20           dimethyl-aminopropyl) carbodiimide (EDAC; Biorad, Richmond, CA.). The aminoethyl derivative was purified by RP-HPLC on a semi-preparative C-4 column using a 5-100% acetonitrile gradient in 0.1% TFA. Eluted material was further purified by S-Sepharose  
25           chromatography. The amino derivative was eluted with 0.1 M HCl, followed by extraction into phenol, and back-extraction into water after the addition of dichloromethane to the phenol phase. Aminoethyl-eVB12 was then recovered from the water phase by  
30           lyophilization.

          The three ANTIDE analogues were conjugated to eVB12 using the carbodiimide, EDAC, and the products (B12-ANT1, B12-ANT2 and B12-ANT3; depending on whether the complex was formed between ANTIDE-1, ANTIDE-2 or  
35           ANTIDE-3, respectively) were purified by chromatography on Sephadex G-25 in 10% acetic acid prior to RP-HPLC on a semi-preparative C-4 column, as described above.









**Conjugation of VB12 to ANTIDE using cross-linker containing a hindered thiol**

Aminoethyl-eVB12 was reacted with 4-succinimidylloxycarbonyl- $\alpha$ -methyl- $\alpha$ -(2-pyridyldithio)toluene (SMPT; Pierce). Iminothiolyated ANTIDE-1 and ANTIDE-3 were prepared as outlined above. The SMPT-derivative of aminoethyl-eVB12 was then reacted with the thiolyated ANTIDE-1 and -3 derivatives as described above, followed by subsequent purification of B12-TSS-ANT1 and B12-TSS-ANT3 by the usual method (Table 2).

**Formation of a non-cleavable thio-ether linkage between VB12 and ANTIDE**

6-(Iodoacetamido)hexylamido-eVB12 in diisopropylethyl amide/dimethyl formamide (DMF/DIEA; 1:20 v/v) was deoxygenated with argon for ten minutes, and a solution of iminothiolyated ANTIDE-1 or -3 in DMF was then added dropwise to the stirred solution. Each reaction mixture was stirred for 30 minutes at room temperature then diluted with water, filtered and the products, B12-SC-ANT1 and B12-SC-ANT-3, were purified by preparative RP-HPLC (Table 2).

Table 2      *Structure of spacer and IF affinity of VB<sub>12</sub>-Antide conjugates*  
*General Structure: eVB<sub>12</sub> - SPACER - Antide*

Conjugate Name	Antide analogue	SPACER	IF affinity
direct	Antide 1	None	N/T
	Antide 3		N/T
anilido	Antide 1		53%
	Antide 3		36%
EGS-derived	Antide 1		30%
	Antide 3		N/T
DSS-derived	Antide 3		N/T
dithio-propionyl	Antide 1		65%
dithio-imino	Antide 3		N/T
hexylthio-imino	Antide 1		82%
	Antide 3		65%
hexyltoluyl-dithio imino	Antide 1		54%
	Antide 3		37%
GGEA	Antide 1		60%
	Antide 3		48%

*N.T. Not tested*



### Conjugation of VB12 to ANTIDE using a transglutaminase cleavable linkage

ANTIDE-1 and -3 were taken up in 5% DIEA/DMF and the *N*-succinimidyl ester derivative of eVB12-GGEA-OMe (eVB12-GlyGlyGluAla) in DMF (100  $\mu$ l) was added. Each reaction mixture was stirred for 30 minutes at room temperature, then diluted with 2% acetic acid, and filtered. The products, B12-TP-ANT1 and B12-TP-ANT3 were purified by RP-HPLC, lyophilized and stored under dessicant before use (Table 2).

### Statistical analysis

Data from replicate individuals (*in vivo* trials), and from replicate experiments (*in vitro* trials) were subjected to one-way analysis of variance, and Tukey HSD post hoc tests were applied for the pairwise comparison of means, which were deemed to be significantly different at a probability level of  $P < 0.05$ .

### Affinity of VB12-ANTIDE conjugates for Intrinsic Factor

All VB12-ANTIDE conjugates tested showed good affinity for Intrinsic Factor (Table 2). In fact, most conjugates had a relative affinity which was higher than that of the eVB12 isomer used to make the conjugate (normally around 35%).

### Example 1

#### Formation of VB12-ANTIDE analogue conjugates using non-cleavable homobi-functional cross-linkers

The direct conjugation between VB12 and ANTIDE-1 and ANTIDE-3 produced conjugates with greatly reduced bioactivity when compared to ANTIDE. The close proximity of VB12 to ANTIDE using this conjugation strategy would sterically interfere with the binding of ANTIDE to the LHRH receptor. In order to reduce the steric effect possibly seen with direct conjugation, non-cleavable linkers were used to produce covalent complexes between aminoethyl"e"VB12 and ANTIDE-1 and 3. Briefly, the ANTIDE-1 and 3 were reacted with a 1.5

molar excess of either Disuccinimidyl suberate (DSS) or Ethylene glycolbis(succinimidylsuccinate) (EGS) for 10 minutes at Room Temperature (RT). Aminoethyl"e"VB12 was then added and the reaction allowed to proceed  
5 overnight. Conjugated material was purified by chromatography on Sephadex G-25 in 10% acetic acid, followed by Reversed Phase HPLC (RP-HPLC). The VB12-Anilido-ANTIDE-1 and ANTIDE-3 conjugates were formed by reaction of aminoethyl"e"VB12 with *p*-aminophenylacetic  
10 acid using EDAC/NHS. The VB12-anilide, was in turn conjugated to ANTIDE-1 and ANTIDE-3 using EDAC/NHS. The conjugated material was purified by G-25 chromatography in 10% acetic acid followed by RP-HPLC.

#### Example 2

#### 15 Formation of VB12-ANTIDE analogue conjugates using spacers

Preparation of conjugates between VB12 and ANTIDE-1 and ANTIDE-3 with either hydrophilic spacers such as EGS (16.1 Å), hydrophobic spacers such as DSS  
20 (11.4 Å), or a shorter bulky anilido group, greatly increased the *in vitro* bio-activity of the conjugates, to levels comparable to that of the parent analogue (Figs 1 & 2). Thus, B12-PA-ANT1, B12-EGS-ANT3 and B12-DSS-ANT3 all had bioactivities comparable to that of  
25 native ANTIDE (Figs 1 & 2). By contrast, the *in vivo* bioactivity of these conjugates when given sc to rats was very low, as evidenced by the failure of these conjugates to cause a significant drop in serum LH levels in castrate rats, even at doses of 100 ug of  
30 analogue. ANTIDE and ANTIDE-3, on the other hand, were active at doses as low as 12.5 ug (Figs 3 & 4). It thus appeared that some factor other than steric hindrance was affecting the *in vivo* activity of these VB12-ANTIDE conjugates (cf. Fig. 1 with 3 and 2 with 4).

### Example 3

#### Formation of VB12-ANTIDE conjugates using thiol-cleavable cross-linkers

Linkage of VB12 to ANTIDE analogues using  
5 extended hydrophilic or hydrophobic spacers produced  
conjugates which had similar *in vitro* activity to  
native ANTIDE. Parenteral administration of these  
conjugates, however, had little effect in lowering  
10 serum LH levels of castrate rats over the range of  
doses tested. These results suggested that although the  
general strategy of increasing the bioactivity of the  
VB12-ANTIDE conjugates by the introduction of extended  
spacers was correct, the conjugate so formed had  
reduced activity *in vivo*, possibly due to the presence  
15 of the VB12 molecule in the conjugate. Conjugates were  
therefore prepared in which the covalent linker  
contained a biodegradable disulfide bond, which would  
be reduced *in vivo*, presumably by glutathione in serum.  
Briefly, aminoethyl"e"VB12 was reacted with N-  
20 succinimidyl 3-(2-pyridyldithio)propionate (SPDP). The  
dithiopyridyl-aminoethyl"e"VB12 (DTP"e"VB12) product  
was purified by RP-HPLC. A free thiol was introduced  
onto ANTIDE-1 by reaction with SPDP. The dithiopyridyl  
group was then reduced with mercapto-ethanol and the  
25 product purified by RP-HPLC. Similarly, a free thiol  
was introduced into ANTIDE-3 by reaction with  
iminothiolane. The thiolated product (SH-HN<sup>+</sup>ANTIDE-3)  
was purified by RP-HPLC. Formation of the disulfide  
linked VB12-ANTIDE-1 and VB12-ANTIDE-2 conjugates was  
30 achieved by reaction of the thiolated ANTIDE derivative  
with DTP"e"VB12 in 2.5% acetic acid for 24 hours. The  
conjugated material was purified by Sephadex G-25  
chromatography, followed by RP-HPLC.

### Example 4

35 Formation of VB12-ANTIDE analogue conjugates using  
biodegradable linkers

In order to reduce the possibility of removal of the VB12-ANTIDE conjugates from the circulation following injection, VB12-ANTIDE conjugates were prepared using a thiol-cleavable, disulfide-containing, spacer. This spacer would potentially allow the cleavage of ANTIDE from VB12 by the low levels of reduced glutathione in serum. Conjugates were therefore prepared between ANTIDE-1 and ANTIDE-3 using a thiol-cleavable spacer (SS) and a hindered thiol group (TSS). The presence of the hindered thiol in the latter spacer should mean that it would be cleaved more slowly upon exposure to serum glutathione. A conjugate containing a non-cleavable thioether linker of similar length (SC) was prepared.

The thiol-cleavable B12-SS-ANT1 and B12-SS-NH<sup>+</sup>ANT3 conjugates were found to have similar *in vitro* bioactivity to the extended spacer conjugates described above (Figs. 1 & 2, Tables 5 & 6). Administration of B12-SS-NH<sup>+</sup>ANT3 parenterally to castrate rats resulted in a similar reduction in serum LH to that seen with the unmodified ANTIDE-3, while B12-SS-ANT1 had a similar bioactivity *in vivo* to native ANTIDE (Table 6). Thus, the use of the thiol-cleavable spacer, rather than a non-cleavable spacer, in the preparation of the VB12-ANTIDE conjugates, while having little effect on the IC50 of the VB12-spacer-ANTIDE conjugates in the *in vitro* assay, greatly increased the potency of these conjugates *in vivo*, to levels similar to ANTIDE (Figs. 3 & 4, Tables 5 & 6). The incorporation of a tolyl-group, or hindered thiol, did not significantly change the *in vitro* or *in vivo* activity of either analogue relative to the respective unhindered thiol analogue. The non-cleavable conjugates formed with the thio-ether linkage were found to have similar activity to the thiol analogue when tested *in vitro*, however, this analogue had a greatly reduced activity when tested *in vivo*.

Table 3. *In vitro* bioassay for blockade of LHRH-stimulated release of LH from rat pituitary cells by ANTIDE, its analogues and VB<sub>12</sub>-conjugates

Analogue	IC <sub>50</sub>	[ <u>n</u> ]
ANTIDE	4.4±0.7 <sup>a</sup>	[20]
ANTIDE-1	2.1±0.3 <sup>a</sup>	[8]
ANTIDE-2	11.7±2.7 <sup>a,b</sup>	[5]
ANTIDE-3	1.2±0.2 <sup>a</sup>	[13]
B12-ANT1	88±21 <sup>c</sup>	[5]
B12-ANT2	105±21 <sup>c</sup>	[4]
B12-ANT3	36±10	[4]

Median inhibitory concentrations (IC<sub>50</sub>) of ANTIDE, ANTIDE analogues and VB12-ANTIDE conjugates for antagonising LHRH-stimulated release of LH from rat anterior pituitary cell cultures during 4 hours. Results for conjugates (mean +/- S.E.) are given as ng of the incorporated Antide analogue/ml (final concentration in culture well) from n independent trials. Results that are not significantly different from each other (i.e. P>0.05) are grouped under a common subscript letter.

*Table 4 Serum LH levels in castrate rats following subcutaneous injection of ANTIDE, ANTIDE analogues or their VB<sub>12</sub>-conjugates*

Analogue	Dose (ug)	Mean LH $\pm$ s.d. (ng/ml)
<u>Expt. 1</u>		
Control		7.35 $\pm$ 1.90 <sup>a*</sup>
ANTIDE	100	0.29 $\pm$ 0.06 <sup>b</sup>
	50	0.24 $\pm$ 0.06 <sup>b</sup>
ANTIDE-1	100	6.37 $\pm$ 1.00 <sup>a</sup>
ANTIDE-2	100	9.51 $\pm$ 3.11 <sup>a</sup>
ANTIDE-3	100	0.27 $\pm$ 0.05 <sup>b</sup>
B12-ANT3	50	10.11 <sup>a</sup>
<u>Expt. 2</u>		
Control		6.69 <sup>a</sup>
ANTIDE	50	0.44 $\pm$ 0.02 <sup>b</sup>
ANTIDE-3	16.7	0.71 $\pm$ 0.2 <sup>b</sup>
B12-ANT1	50	5.44 $\pm$ 1.4 <sup>a</sup>
B12-ANT2	50	5.56 $\pm$ 0.97 <sup>a</sup>
B12-ANT3	50	5.51 $\pm$ 0.22 <sup>a</sup>

\* A common letter signifies that the mean values do not significantly differ from one another (P>0.05)

Table 5. *In vitro* bioassay for blockade of LHRH-stimulated release of LH from rat pituitary cells by ANTIDE, its analogues and disulfide linked VB<sub>12</sub>-conjugates.

Analogue	IC50*	[n]
ANTIDE	4.4±0.7 <sup>a</sup>	[20]
ANTIDE-1	2.1±0.3 <sup>a</sup>	[8]
B12-SS-ANT1	10.4±4.0 <sup>b</sup>	[3]
ANTIDE-3	1.2±0.2 <sup>c</sup>	[15]
HS-NH <sup>+</sup> ANT3	0.72±0.11 <sup>a,c</sup>	[3]
B12-SS-NH <sup>+</sup> ANT3	7.4±0.8 <sup>a,b</sup>	[3]

\* (See legend to Table 3 for further details).

*Table 6      Reduction in serum LH level in castrate rats following subcutaneous injection of ANTIDE, ANTIDE analogues, or their disulfide linked VB<sub>12</sub>-conjugates*

Analyse	Dose (ug) of Analyse	Serum LH, % of matching control* (mean $\pm$ s.d.)
ANTIDE	25.0	4 $\pm$ 0.70    [#1]
	12.5	15.3 $\pm$ 5.40    [#3]
	6.3	80.6 $\pm$ 43.8    [#3]
	3.1	100.2 $\pm$ 36.8    [#3]
ANTIDE-1	100	87.6 $\pm$ 13.6    [#1]
B12-SS-ANT1	25.0	19.8 $\pm$ 4.80    [#5]
	12.5	34.4 $\pm$ 14.7    [#5]
	6.3	73.9 $\pm$ 25.4    [#6]
ANTIDE-3	100.0	3.7 $\pm$ 0.70    [#1]
	25	18.0 $\pm$ 5.50    [#5]
	12.5	43.0 $\pm$ 12.8    [#3]
	6.3	116.1 $\pm$ 42.4    [#3]
HS-NH <sup>+</sup> ANT3	50.0	28.2 $\pm$ 11.4    [#5]
	25.0	77.3 $\pm$ 17.2    [#5]
	12.5	112.8 $\pm$ 20.9    [#5]
B12-SS-NH <sup>+</sup> ANT3	50.0	16.5 $\pm$ 5.10    [#5]
	25.0	23.8 $\pm$ 9.20    [#5]
	12.5	63.8 $\pm$ 16.5    [#6]
	6.3	78.8 $\pm$ 24.4    [#6]



**Example 5**

**Formation of VB12-ANTIDE analogue conjugates using a transglutaminase cleavable linker.**

Conjugates formed between VB12 and ANTIDE-1 and  
5 ANTIDE-3 using a transglutaminase cleavable  $\gamma$ -glutamyl- $\epsilon$ -lysine bond, (B12-TP-ANT1 and B12-TP-ANT3) were found to have poor activity *in vitro*, and negligible activity *in vivo* at the doses tested, suggesting that this bond was not cleaved *in vitro* or *in vivo*. The stereo-  
10 specificity may have inhibited the cleavage of the ANTIDE-1 conjugates (due to the presence of the D-lysine group), while steric hindrance may have prevented the transglutaminase enzyme from cleaving both the ANTIDE-1 and ANTIDE-3 conjugates.

**15 Example 6**

**Formation of VB12-ANTIDE-Analogue Lysyl-Polyglutamate conjugates using thiol-cleavable cross-linkers**

It is desirable to increase the amount of drug, or its analogues, which can be taken up by the VB12-  
20 transport system, by the linkage of multiple copies of the drug to a polymeric backbone, to which is conjugated one or more VB12 molecules. A process is described for the formation of such a VB12-drug-polymer complexes.

**25 a) Formation of DTP-Lysyl-polyglutamate**

Poly-glutamate (100 mg) (MW 64,600-70,000; Sigma), was reacted with EDAC (100 mg) and NHS (50 mg in acetone) for 10 minutes at RT. Lysine (400 mg in 4 ml 1%NaHCO<sub>3</sub>) was added and allowed to react overnight  
30 (O/N). The product, Lysyl-Polyglutamate (LPG), was purified by extensive dialysis against DW, and then lyophilized.

The dithiopyridyl derivative of LPT, was obtained by reacting Dithiopyridyl-propionic acid (50 mg) with  
35 O-(N-Succinimidyl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TSTU) (100 mg) and N-Ethyldiisopropyl amine (100 mg; DIEA) in DMF for 1

hour. The succinimidyl ester so formed was added directly to 100 mg LPG dissolved in 4 ml 2% NaHCO<sub>3</sub>. The reaction was allowed to proceed O/N after which the product (DTP-LPG) was purified by exhaustive dialysis against DW, and then lyophilized.

b) *Preparation of Iminothiolated ANTIDE-1*

ANTIDE-1 (20 mg) was dissolved in 300 µl 5% DIEA in DMF containing 5 mg EDTA plus 5 mg DTT, which had been degassed under argon. Iminothiolane (20 mg in 50 µl DIEA/DMF plus 50 µl Borate buffer, 100 mM pH 8.2) was added and reacted for 60 minutes before purification by RP-HPLC and lyophilization.

c) *Preparation of Iminothiolated aminoethyl-VB<sub>12</sub>*

Aminoethyl-VB<sub>12</sub> (20 mg) was dissolved in 300 µl 5% DIEA in DMF containing 5 mg EDTA plus 5 mg DTT, which had been degassed under argon. Iminothiolane (20 mg in 50 µl DIEA/DMF plus 50 µl Borate buffer, 100 mM pH 8.2) was added and reacted for 60 minutes before purification by RP-HPLC and lyophilization.

d) *Formation of VB<sub>12</sub>-ANTIDE-Lysyl-Polyglutamate*

DTP-LPG was dissolved at 20 mg/ml in DW. Iminothiolated-aminoethyl-VB<sub>12</sub> was dissolved at 5 mg/ml in DW and added to the DTP-LPG (1:20 w/w) and allowed to react for 20 min at RT before the addition of iminothiolated-ANTIDE-1 (50 mg/ml in DW) dropwise with stirring. The reaction mixture was kept at pH 6.5-7.0 by the addition of Tris.HCL pH 7.0 and NaAcetate, pH 5.5. The reaction proceeded overnight after which the product was purified by dialysis and then lyophilized. The composition of the product was determined by amino acid analysis and found to contain 1 : 5 : 21, ANTIDE-1 : Lysine : Glutamate, or roughly 25% by weight ANTIDE.

The development of an LHRH antagonist suitable for use in humans for the treatment of hormone dependent mammary and prostate cancer has in the past been limited by the low potency of these molecules, their poor solubility in physiological buffers, and their poor oral bioavailability. In an attempt to resolve these problems, the present inventor has examined the possibility of combining the relatively high bioactivity of the potent LHRH antagonist, ANTIDE, with the high oral bioavailability of vitamin B12. ANTIDE is particularly suitable for oral delivery, since it is composed of many unnatural amino acids, and is resistant to the action of trypsin or chymotrypsin. ANTIDE, however, must be modified in such a way as to provide a suitable site for conjugation to VB12, as well as retaining full biological potency. ANTIDE normally possesses modified lysine residues at positions 5, 6 and 8 viz:- Lys(Nic), D-Lys(Nic) and Lys(iPr), respectively. The removal of these modifications at either of the three sites provides a free amine suitable for conjugation to VB12. Two derivatives of ANTIDE, D-Lys<sup>6</sup>-ANTIDE (ANTIDE-1) and Lys<sup>8</sup>-ANTIDE (ANTIDE-3) suitable for conjugation to VB12 were synthesized, and found to have high potency in the *in vitro* pituitary cell assay. ANTIDE-3 was also found to be highly potent when administered s.c. to rats, although ANTIDE-1, was found to have negligible antagonist activity, possibly due to an alteration in the rate of its clearance from serum. The *in vitro* and *in vivo* potency of either analogue was, however, greatly reduced by direct conjugation to VB12 (Tables 3 & 4), presumably due to steric interference with binding of the conjugate to the pituitary receptor for LHRH by the bulky VB12 molecule in close proximity to ANTIDE.

In order to reduce the steric effect seen with direct conjugation of VB12 to ANTIDE-1 and ANTIDE-3,

non-cleavable linkers were used to produce covalent complexes between aminoethyl-eVB12 and ANTIDE-1 and -3, thereby separating the bulky VB12 group from the ANTIDE analogue. VB12-ANTIDE-1 conjugates formed with  
5 either an anilido (PA) or an EGS spacer, or VB12-ANTIDE-3 conjugates formed with an anilido, EGS- or DSS-derived spacer, all produced conjugates with *in vitro* activity similar to that of native ANTIDE (Fig 1), suggesting that placing the VB12 group at some  
10 distance from the ANTIDE restored the affinity of the ANTIDE for its receptor. These conjugates had very low activities *in vivo* (Fig. 2), possibly due to the removal of the conjugates from the circulation by VB12-binding proteins, such as Transcobalamin I or  
15 Transcobalamin II, which would greatly reduce their *in vivo* potency.

In an attempt to reduce the possibility of steric hindrance produced by the presence of the VB12 in the conjugates, or reduced systemic bioavailability of the  
20 conjugates as a result of clearance by VB12-binding proteins, VB12-ANTIDE conjugates were prepared using linkers which were potentially biodegradable *in vivo*. A disulfide linkage was initially chosen, because this linkage is most readily cleaved *in vivo*, and is found  
25 in biological conjugates such as the A-SS-B subunit toxins, tetanus toxin and diphtheria toxin (Sandvig & Olsens, 1981; Matsuda & Yoneda, 1974). Disulfide bonds have been shown to be cleaved by the low levels of glutathione found in serum, or in the cytoplasm of  
30 cells (Anderson & Meister, 1980).

The disulfide-linked VB12-ANTIDE-1 and VB12-ANTIDE-3 conjugates were found to have a slightly reduced *in vitro* potency in comparison to ANTIDE, or the Anilido- and EGS-linked VB12-ANTIDE-1 and ANTIDE-3  
35 conjugates. In contrast, the disulfide linked conjugates showed a dramatic increase in their *in vivo* potencies when compared to the non-cleavable, extended

spacer conjugates (Figs. 3 & 4, Tables 5 & 6). For the B12-SS-NH<sup>+</sup>-ANT3 conjugate, the *in vivo* bio-activity slightly exceeded that of the HS-NH<sup>+</sup>-ANT3 analogue. A spacer of similar length containing a non-cleavable thioether bond rather than the disulfide bond showed greatly reduced *in vivo* activity, suggesting that the increased activity of the disulfide linked conjugate was due to its ability to be cleaved *in vivo*. All conjugates described above, whether they were directly linked to VB12 or were linked by a cleavable or non-cleavable spacer were found to be significantly more soluble in physiological buffers such as PBS or saline, than the parent ANTIDE molecule (results not shown).

Conjugates between VB12 and ANTIDE-1 and -3 were formed with a spacer containing a  $\gamma$ -glutamyl- $\epsilon$ -lysine bond, which could potentially be cleaved by serum transglutaminases. Despite the high *in vitro* activity of these conjugates, their *in vivo* activity was very low, suggesting that the spacer was not cleaved *in vivo*.

The affinity of the VB12-ANTIDE-1 and -3 conjugates for IF was found to be equal or greater than the affinity of the eVB12 isomer from which the conjugates were made. A similar observation has been made for other eVB12 isomer conjugates (Russell-Jones, 1994)

These studies demonstrate that it is possible to link together two molecules with disparate biological activities and to substantially maintain the biological activity of each individual component. During such conjugation, care must be taken to maintain the receptor-ligand binding activity of both molecules, as well as preserving the bio-availability of the pharmacological agent. Thus, although it was possible to make several complexes between VB12 and ANTIDE using spacers while showing high *in vitro* activity in the pituitary cell assay, and with high relative affinity

for IF, the *in vivo* bioactivity of these conjugates was greatly reduced. In this case, an alternative approach using biodegradable spacers to join the two molecules proved successful.

- 5           It will be appreciated by persons skilled in the art that numerous variations and/or modifications may be made to the invention as shown in the specific embodiments without departing from the spirit or scope of the invention as broadly described. The present  
10           embodiments are, therefore, to be considered in all respects as illustrative and not restrictive.

## REFERENCES

1. Schally, A. B., Ariumara, A., Kastin, A. J., et al., 1971 Science 173, 1036
2. Loumay, E., Naor, Z., and Catt, K. J. 1982 Endocrinol., 111, 730
3. Perrin, M. H., Rivier, J. E., and Vale, W. W. 1980 Endocrinol., 102, 1289
4. Eisenberger, M., and Abrams, J., 1988., Drugs Today, 24, 241
5. Crawford, D. D., and Davis, M. A., in "Endocrine Therapies in Breast and Prostate Cancer." (Ed) C. K. Osborne, Boston p.25.
6. Karten, M. J., and Rivier, J. E. 1986 Endocrin. Rev., 7, 44
7. Schmidt, F., et al., 1984 Contraception, 29, 283
8. Phillips, A., et al., 1988 Life Sciences 43, 883.
9. Ljunggquist, A., et al., 1988 P.N.A.S. 85, 8236
10. Leal, J. A., et al., 1988 J. Clin. Endocrinol. Metab., 67, 1325
11. Habenicht, U.-F., Schneider, M. R., Fathy El Etreby, M. 1990 J. Steroid. Biochem. Molec. Biol., 37, 397.
12. McLachlan, T. I., Healy, D. L., Burger, H. G., 1986 Brit. J. Obstet. Gynaec., 93, 431.
13. Farnworth, P. G., Robertson, D. M., de Kretser, D. M. and Burger, H. G. 1988 Effects of 31 kilodalton bovine inhibin on follicle-stimulating hormone and luteinizing hormone in rat pituitary cells *in vitro*: actions under basal conditions. Endocrinol., 122, 207-313.
14. Russell-Jones, G. J. (1994) Oral delivery of Therapeutic Proteins and peptides by the vitamin B12 uptake system. In "Peptide-based Drug Design: Controlling Transport and Metabolism." Ed. M. Taylor and G. Amidon, ACS Publications.

15. Sandvig, K. & Olsens, S. (1981) Rapid entry of nicked Diptheria toxin into cells at low pH. Characterization of the entry process and effects of low pH on the toxin molecule. *J. Biol. Chem.*, 256, 9068-9076.
16. Matsuda, M. & Yoneda, M (1974) Dissociation of tetanus neurotoxin into two polypeptide fragments. *Biochem. Biophys. Res. Com.*, 57, 1257-1262.
17. Anderson, M. E., & Meister, A. (1980) Dynamic state of glutathione in blood plasma. *J. Biol. Chem.*, 255, 9530-9533.



## CLAIMS

1. A LHRH antagonist selected from the group consisting of N-Ac-D-Nal(2), D-Phe (pCl), D-Pal(3), Ser, Lys (Nic), D-Lys, Leu, Lys(iPr), Pro, D-Ala-NH<sub>2</sub>  
5 (D-Lys<sub>6</sub> ANTIDE or ANTIDE-1); N-Ac-D-Nal(2), D-Phe (pCl), D-Pal(3), Ser, Lys, D-Lys(Nic), Leu, Lys(iPr), Pro, D-Ala-NH<sub>2</sub> (Lys<sub>5</sub> ANTIDE or ANTIDE-2) and N-Ac-D-Nal(2), D-Phe (pCl), D-Pal(3), Ser, Lys (Nic), D-Lys(Nic), Leu, Lys, Pro, D-Ala-NH<sub>2</sub> (Lys<sub>8</sub>ANTIDE or  
10 ANTIDE-3).
2. A complex for use in chemical castration of a subject and/or in treatment of gonadotrophin dependent disorders the complex comprising an LHRH antagonist linked to a carrier being vitamin B12 or an analogue  
15 thereof via a linker group, the linker group being characterised in that the LHRH antagonist is cleaved from a carrier *in vivo*.
3. A complex as claimed in claim 2 wherein the LHRH antagonist is selected from the group consisting of  
20 ANTIDE-1, ANTIDE-2 and ANTIDE-3.
4. A complex as claimed in claim 3 wherein the LHRH antagonist is ANTIDE-3.
5. A complex as claimed in any one of claims 2 to 4 wherein more than one LHRH antagonist is linked to the  
25 carrier.
6. A complex as claimed in any one of claims 2 to 5 in which the carrier is selected from the group consisting of cyanocobalamin, aquocobalamin, adenosylcobalamin, methylcobalamin, hydroxycobalamin,  
30 cyanocobalamin, carbanalide, 5-o-methylbenzylcobalmin, the desdimethyl, monoethylamide and the methylamide analogues all of the above, analogues and homologues of cobamamide, coenzyme B12, 5'-deoxyadenosylcobalamine, chlorocobalamin, sulfitocobalamin, nitrocobalamin,  
35 thiocyanatocobalamin, benzimidazole derivatives, adenosylcyanocobalmin, cobalamin lactone, cobalamin lactam and the anilide, ethylamide, propionamide,

monocarboxylic and dicarboxylic acid derivatives of vitamin B12 or its analogues.

7. A complex as claimed in any one of claims 2 to 6  
5 in which the LHRH antagonist is linked to the carrier  
via a thiol cleavable cross-linker.

8. A complex as claimed in claim 6 in which the LHRH  
antagonist is linked to the carrier by a disulphide  
linkage.

9. A method for chemical castration and/or treatment  
10 of gonadotrophin dependent disorders of a subject  
comprising administering to the subject an effective  
amount of the complex as claimed in any one of claims 2  
to 8.

1/4




Fig. 1

2/4




Fig. 2

3/4




Fig. 3

4/4





Fig. 4

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU 94/00262

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> Int. Cl. <sup>5</sup> C07K 007/20 A61K 047/24, 37/43  According to International Patent Classification (IPC) or to both national classification and IPC					
<b>B. FIELDS SEARCHED</b>  Minimum documentation searched (classification system followed by classification symbols) IPC:C07K 007/20, 7/06, A61K 47/24, 37/43  Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched AU:IPC as above  Electronic data base consulted during the international search (name of data base, and where practicable, search terms used) DERWENT: "LHRH" or "luteinising hormone releasing hormone", "antagonist" and "carrier" or "complex" Chemical Abstracts protein sequence search					
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>					
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to Claim N .			
X	AU-B-65289/86 (587658) (Biotechnology Australia Pty Ltd.) 5 May 1987 (05.05.87) See claims 1-6	2-9			
X	AU-B-32637/89 (623385) (Sandoz Ltd.) 12 October 1989 (12.10.89) See claims.	1			
A	AU-A-80593/91 (The Salk Institute for Biological Studies) 7 January 1992 (07.01.92)				
<div style="display: flex; justify-content: space-between;"> <div> <input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C.         </div> <div> <input checked="" type="checkbox"/> See patent family annex.         </div> </div>					
<table style="width: 100%; border: none;"> <tr> <td style="width: 33%; vertical-align: top;">           * Special categories of cited documents :             "A" document defining the general state of the art which is not considered to be of particular relevance            "E" earlier document but published on or after the international filing date            "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)            "O" document referring to an oral disclosure, use, exhibition or other means            "P" document published prior to the international filing date but later than the priority date claimed         </td> <td style="width: 33%; vertical-align: top;">           "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention            "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone            "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art            "&amp;" document member of the same patent family         </td> <td style="width: 33%;"></td> </tr> </table>			* Special categories of cited documents :  "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family	
* Special categories of cited documents :  "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family				
Date of the actual completion of the international search 29 August 1994 (29.08.94)	Date of mailing of the international search report <b>07 SEP 1994 ( 07.09.94 )</b>				
Name and mailing address of the ISA/AU  AUSTRALIAN INDUSTRIAL PROPERTY ORGANISATION PO BOX 200 WODEN ACT 2606 AUSTRALIA  Facsimile No. 06 2853929	Authorized officer   <b>T. SUMMERS</b>  Telephone No. (06) 2832291				

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU 94/00262

Category*	Citation of document, with indication, where appropriate of the relevant passages	Relevant to Claim No.
A	Ljungqvist, A. et al "Antagonists of LHRH Superior to Antide; ....." Tetrahedron Vol. 46 No. 9 pp 3297-3304, 1990	
A	Ljungqvist, A. et al "Antide and related antagonists of luteinizing ....." Proc. Nat. Proc. Nat. Acad. Sci. USA Vol. 85 pp 8236-8240 November 1988	
A	Phillips, A. et al "Evaluation of the anaphylactoid activity of a new LHRH antagonist". Life Sciences Vol. 43 No. 11 pp 883-888 1988	



**INTERNATIONAL SEARCH REPORT**  
Information on patent family member

International application No.  
**PCT/AU 94/00262**

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report		Patent Family Member	
AU-B 65289/86	KR 940357B1	NZ 217821	
CN 860107590	ZA 8607703-A	DK2925/87-A1	
EP 220030-B1	AT- 64584-E	GR3002160-T3	
IL 80264-A1	DE- 3679882-CO	JP63501015-T2	ES 2051690-T3
	CA 1330791-A1	IN 165029-A	WO 8702251-A1
AU-B 32637/89	NL 8900864-A	ES 2105645-A	
AU-A 33628/89	LU 87495-A	ZA89052570-A	
WO-A 8909786	DK 8901636-A	BE1004357-A4	
GB 2218102-A	JP 01305099-A	IT 1235894-B	
GB 2247682-A	SE 8901241-A	DE 3990350-A	
	CH 682152-A5		
END OF ANNEX			

**THIS PAGE BLANK (USPTO)**